CPC(HEP & NP), 2009, **33**(X): 1—10 **Chinese Physics C** Vol. 33, No. X, Xxx, 2009

arXiv:1111.3533v1 [hep-ph] 15 Nov 2011

$X(1835)$, $X(2120)$, $X(2370)$ and $\eta(1760)$ in chiral quark model *

Shaona Chen1) Jialun Ping[1,2;1)]

[1] School of Physical Science and Technology, Nanjing Normal University, Nanjing 210097, China

[2] School of Physics and Microelectronics, Shandong University, Jinan 250100, China

Abstract: By analyzing the meson spectra obtained in the constituent quark model, we find that the pseudoscalar mesons $\eta'(2^1S_0)$, $\eta(4^1S_0)$, $\eta'(3^1S_0)$ and $\eta'(4^1S_0)$ are the possible candidates of $\eta(1760)$, $X(1835)$, $X(2120)$ and $X(2370)$. The strong decay widths of these pseudoscalars to all the possible decay modes are calculated within the framework of the $^3P_0$ model. Although the total width of $\eta'(2^1S_0)$ is compatible with the experimental value of BES for $\eta(1760)$, the partial decay width to $\omega\omega$ is too small, which is not consistent with the result of BES. If the state $X(1835)$ is interpreted as $\eta(4^1S_0)$, the total decay width is compatible with the experimental data, and the main decay modes will be $\pi a_0(980)$ and $\pi a_0(1450)$, which needs to be checked. The assignment of of $X(2120)$ and $X(2370)$ to $\eta'(3^1S_0)$ and $\eta'(4^1S_0)$ is also disfavored in the present calculation because of the incompatibility of the decay widths.

Key words: X(1835), X(2120), X(2370), $\eta(1760)$, chiral quark model, pseudoscalar meson

PACS: 12.39.Jh, 13.25.Jx, 14.40.Aq

1 Introduction

In 2005, the BES Collaboration observed a narrow peak in the $\eta'\pi^+\pi^-$ invariant mass spectrum in the process $J/\psi \to \eta'\pi^+\pi^-$ with a statistic significance of 7.7σ. Fitting with Breit-Wigner function yields mass and width [1]

$$\begin{aligned} M &= 1833.7 \pm 6.1(stat) \pm 2.7(syst)\ \mathrm{MeV/c^2} \\ \Gamma &= 67.7 \pm 20.3(stat) \pm 7.7(syst)\ \mathrm{MeV/c^2}, \end{aligned}$$

and the product branching fraction

$$\begin{aligned} &B(J/\psi \to \gamma X(1835))B(X(1835) \to \pi^+\pi^-\eta') \\ &\quad = (2.2 \pm 0.4(stat) \pm 0.4(syst)) \times 10^{-4}. \end{aligned}$$

BES-III confirmed it in the same process with statistical significance larger than 20σ. The fitted mass and width are $M = 1836.5\pm3.0(\mathrm{stat})^{+5.6}_{-2.1}(\mathrm{syst})$ MeV/c^2, Γ=190 $\pm9(\mathrm{stat})^{+38}_{-36}(\mathrm{syst})$ MeV/c^2. Meanwhile, another two new resonances, $X(2120)$ and $X(2370)$, are also observed in the same process with the statistical significance larger than 7.2σ and 6.4σ, respectively. The fitted masses and widths are [2]

$$\begin{aligned} M &= 2122.4 \pm 6.7(stat)^{+4.7}_{-2.7}(syst)\ \mathrm{MeV/c^2}, \\ M &= 2376.3 \pm 8.7(stat)^{+3.2}_{-4.3}(syst)\ \mathrm{MeV/c^2}, \end{aligned}$$

and

$$\begin{aligned} \Gamma &= 83 \pm 16(stat)^{+31}_{-17}(syst)\ \mathrm{MeV/c^2}, \\ \Gamma &= 83 \pm 17(stat)^{+44}_{-6}(syst)\ \mathrm{MeV/c^2}, \end{aligned}$$

respectively. $\eta(1760)$, which its nature is in controversial, was first reported by Mark III collaboration in the J/ψ radiative decays to $\omega\omega$ [3] and $\rho\rho$ [4]. And DM2 collaboration observed a large bump peaked at 1.77 GeV/c^2 in $\omega\omega$ invariant mass distribution in the process of $J/\psi \to \gamma\omega\omega$ ($\omega \to \pi^+\pi^-\pi^0$) [5] and the study of the decays $J/\psi \to \gamma\pi^+\pi^-\pi^+\pi^-$ and $J/\psi \to \gamma\pi^+\pi^-\pi^0\pi^0$ showed that both decays have a large $\rho\rho$ dynamics [6]. The fitted mass and width are M = 1760±11 MeV, $\Gamma = 60 \pm 16$ MeV. Recently BES collaboration reported its results on the decays $J/\psi \to \gamma\omega\omega$, $\omega \to \pi^+\pi^-\pi_0$ [7]. The mass and width turn to be $M = 1744 \pm 10(\mathrm{stat})\pm15$ MeV, Γ=$244^{+24}_{-21} \pm 25$ Mev.

Many works have been devoted to the underlying structures of $X(1835)$ and $\eta(1760)$ [8]. For $X(1835)$, Some interpret it as a $p\bar{p}$ bound state [9–12]. By calculating the mesonic decays of a baryonium resonance, Ding *et al.* claimed that the $p\bar{p}$ bound state favors the decay channel $X \to \eta 4\pi$ over $X \to \eta 3\pi$ [9]. In fact, it is just this work that stimulates the obser-

---

* Supported by National Natural Science Foundation of China (11035006 and 11175088)

1) E-mail: jlping@njnu.edu.cn (Corresponding author)

©2009 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institut

vation of $J/\psi \to \eta' \pi^+ \pi^-$ process in BES experiments. Using a semi-phenomenological potential model that can describe all the $N\bar{N}$ scattering data, Dedonder *et al.* found a broad spin-isospin singlets, S-wave quasi-bound state of $N\bar{N}$, which can be used to explain the observed peak by BES [10]. Z. G. Wang and S. L. Wang also calculated the mass of $X(1835)$, which as a baryonium in the framework of QCD sum rule and obtained a consistent result with experimental data. The large-N_c QCD is also applied to study the state $X(1835)$ as a baryonium [11]. Interpretation of $X(1835)$ as a glueball or a glueball mixed with pseudoscalar meson or baryonium is also proposed by using QCD sum rule [13–16]. Apart from these explanation for $X(1835)$ as an exotic state, the conventional $q\bar{q}$ picture of $X(1835)$ is also proposed. Huang and Zhu studied the behavior of $X(1835)$ and thought that it can be taken as the second radial excitation of $\eta'(958)$, in the effective Lagrangian approach [17]. The two-body decays of $X(1835)$ as 3^1S_0 are also calculated by quark-pair creation (QPC, or $^3P_0$) model [18], the results show that the decay width is sensitive to the mixing angle of two states $X_n = (u\bar{u} + d\bar{d})/\sqrt(2)$ and $X_s = s\bar{s}$. Recently J. S. Yu *et al.* systematically studied the two-body strong decays and double pion decays of η-family and assigned the $X(1835)$ to the second radial excitation of $\eta'(958)$, $X(2120$ and $X(2370)$ to the third and fourth radial excitation of $\eta(548)/\eta'(958)$, respectively [19]. For $\eta(1760)$, J. Vijande *et al.* assigned it to be 2^1S_0 state of $s\bar{s}$ in the chiral quark model [20]. The assignment of $\eta(1760)$ to the second radial excitation of $\eta(548)$ is also proposed by J. S. Yu *et al.*[19]. Li and Page suggested it to be a gluonic meson [21]. Glueball mixed with $q\bar{q}$ picture of $\eta(1760)$ was also suggested by N. Wu *et al.* [22]. Stimulated by these experimental and theoretical work, we shall study whether $\eta(1760)$, $X(1835)$, $X(2120)$ and $X(2370)$ can be described in the simplest system-$q\bar{q}$ system.

In this work, the pseudoscalar meson spectrum is determined by the chiral quark model, the mixing angle between X_n and X_s is fixed through the system dynamics. Based on the mass spectrum, the possible candidates of $X(1835)$, $X(2120)$, $X(2370)$ and $\eta(1760)$ are assigned. Then the strong decay widths of the states are calculated in the framework of $^3P_0$ model. to see the assignment is reasonable or not. The paper is organized as follows: a brief review of $^3P_0$ model is given in section 2. The chiral quark model is introduced and meson spectrum and wave function scale parameter β of the involved mesons are obtained in section 3. The numerical result of the strong decay are shown in section 4. The last section is a summary.

2 Review of $^3P_0$ model of meson decay

The $^3P_0$ model also known as the Quark-Pair Creation (QPC) model, applied to the decay of meson A to meson $B+C$ was first proposed by Micu [23], and then developed by Le Yaouanc, Ackleh, Roberts et al [24–26]. The $^3P_0$ model assumes that there is a pair of quark and antiquark created in vacuum. The quantum number of the pair of quark and antiquark is $J^{pc} = 0^{++}$. Since vacuum is colorless and flavorless, so color and flavor singlet should be satisfied. The created pair recombines with the quark-antiquark pair in initial meson and form two mesons in the final state in two possible ways, which is shown in Fig.1.



Fig. 1. The two possible diagrams contributing to $A \to B+C$ in the $^3P_0$ model.

In the non-relativistic limit, the transition operator T takes form as

$$T = -3\,\gamma \sum_m \langle 1m1-m|00\rangle \int d\mathbf{p}_3 d\mathbf{p}_4 \delta^3(\mathbf{p}_3+\mathbf{p}_4) \times \mathcal{Y}_1^m\left(\frac{\mathbf{p}_3-\mathbf{p}_4}{2}\right) \chi^{34}_{1-m} \phi_0^{34} \omega_0^{34} b_3^\dagger(\mathbf{p}_3) d_4^\dagger(\mathbf{p}_4), \quad (1)$$

where γ, which is a dimensionless parameter, represents the strength of the quark-antiquark pair creation from the vacuum and can be obtained by fitting the experimental data. $\mathbf{p_3}$ and $\mathbf{p_4}$ denote the momenta of the created quark and antiquark respectively. $\mathcal{Y}_l^m(\mathbf{p}) = |p|^l Y_l^m(\theta_p, \phi_p)$ is the l-th solid harmonic polynomial that gives the momentum-space distribution of the created quark-antiquark pair. χ^{34}_{1-m} reflects triplet state of spin. $\phi_0^{34} = (u\bar{u}+d\bar{d}+s\bar{s})/\sqrt{3}$ and $\omega_0^{34} = (r\bar{r}+g\bar{g}+b\bar{b})/\sqrt{3}$ correspond to flavor and color singlets, respectively. $b_3^\dagger(\mathbf{p}_3) d_4^\dagger(\mathbf{p}_4)$ are the creation operators of the quark and antiquark, respectively.

To depict the meson state, we define

$$\begin{aligned}&|A(n_A{}^{2S_A+1}L_{A\,J_AM_{J_A}})(\mathbf{P}_A)\rangle \equiv\\ &\sqrt{2E_A}\sum_{M_{L_A},M_{S_A}}\langle L_AM_{L_A}S_AM_{S_A}|J_AM_{J_A}\rangle\\ &\int d\mathbf{p}_A\chi^{12}_{S_AM_{S_A}}\phi^{12}_A\omega^{12}_A\left|q_1\left(\frac{m1}{m1+m2}\mathbf{P}_A+p_A\right)\right.\\ &\left.\bar{q}_2\left(\frac{m2}{m1+m2}\mathbf{P}_A+p_A\right)\right\rangle, \qquad (2)\end{aligned}$$

The wave function is normalized to

$$\begin{aligned}&\left\langle A(n_A{}^{2S_A+1}L_{A\,J_AM_{J_A}})(\mathbf{P}_A)\right| \qquad (3)\\ &\left.A(n_A{}^{2S_A+1}L_{A\,J_AM_{J_A}})(\mathbf{P}'_A)\right\rangle = 2E_A\delta^3(\mathbf{P}_A-\mathbf{P}'_A).\end{aligned}$$

where $\chi^{12}_{S_AM_{S_A}}, \phi^{12}_A, \omega^{12}_A$ represent the spin, flavor and color wave function respectively; $\mathbf{P_A}$ is the CM momentum of meson A, and $\mathbf{p_A} = (m_2\mathbf{p_1} - m_1\mathbf{p_2})/(m_1+m_2)$ is the relative momentum of $q\bar{q}$ pair. n_A is the radial quantum number; $|L_A, M_{L_A}\rangle, |S_A, M_{S_A}\rangle, |J_A, M_{J_A}\rangle$ are the quantum number of orbit angular momentum between $q\bar{q}$ pair in meson A, the total spin of the pair and the total angular momentum, respectively; $\langle L_AM_{L_A}S_AM_{S_A}|J_AM_{J_A}\rangle$ denotes a Clebsch-Gordan coefficient, E_A is the total energy of the meson.

To describe a strong decay process of $A\to B+C$, the S-matrix is written out as

$$\langle BC|S|A\rangle = I - 2\pi i\delta(E_A-E_B-E_C)\langle BC|T|A\rangle, \quad (4)$$

and then

$$\langle BC|T|A\rangle = \delta^3(\mathbf{P}_A-\mathbf{P}_B-\mathbf{P}_C)\mathcal{M}^{M_{J_A}M_{J_B}M_{J_C}}, \quad (5)$$

where $\mathcal{M}^{M_{J_A}M_{J_B}M_{J_C}}$ is the helicity amplitude of $A\to B+C$. Taking the center of the mass frame of the meson A: $\mathbf{P_A}=0$. One can obtained $\mathcal{M}^{M_{J_A}M_{J_B}M_{J_C}}$ for decay process in terms of overlap integrals,

$$\begin{aligned}\mathcal{M}^{M_{J_A}M_{J_B}M_{J_C}} &= 3\gamma\sum_{\{M\}}\langle L_AM_{L_A}S_AM_{S_A}|J_AM_{J_A}\rangle\langle L_BM_{L_B}S_BM_{S_B}|J_BM_{J_B}\rangle\langle L_CM_{L_C}S_CM_{S_C}|J_CM_{J_C}\rangle\\ &\times \langle 1m1-m|00\rangle\langle\chi^{14}_{S_BM_{S_B}}\chi^{32}_{S_CM_{S_C}}|\chi^{12}_{S_AM_{S_A}}\chi^{34}_{1-m}\rangle[\langle\omega^{14}_B\omega^{32}_C|\omega^{12}_A\omega^{34}_0\rangle\langle\phi^{14}_B\phi^{32}_C|\phi^{12}_A\phi^{34}_0\rangle\\ &\times \mathcal{I}^{M_{L_A},m}_{M_{L_B},M_{L_C}}(\mathbf{P},m_1,m_2,m_3) + (-1)^{1+S_A+S_B+S_C}\langle\omega^{32}_B\omega^{14}_C|\omega^{12}_A\omega^{34}_0\rangle\langle\phi^{32}_B\phi^{14}_C|\phi^{12}_A\phi^{34}_0\rangle\\ &\times \mathcal{I}^{M_{L_A},m}_{M_{L_B},M_{L_C}}(-\mathbf{P},m_2,m_1,m_3)] \qquad (6)\end{aligned}$$

where $\{M\} = M_{L_A}, M_{S_A}, M_{L_B}, M_{S_B}, M_{L_C}, M_{S_C}, m$, The momentum space integral $\mathcal{I}^{M_{L_A},m}_{M_{L_B},M_{L_C}}(\mathbf{P},m_1,m_2,m_3)$ and $\mathcal{I}^{M_{L_A},m}_{M_{L_B},M_{L_C}}(-\mathbf{P},m_2,m_1,m_3)$ are given by

$$\begin{aligned}\mathcal{I}^{M_{L_A},m}_{M_{L_B},M_{L_C}}(\mathbf{P},m_1,m_2,m_3) &= \sqrt{8E_AE_BE_C}\int d\mathbf{p}\,\psi^*_{n_BL_BM_{L_B}}(\tfrac{m_3}{m_1+m_3}\mathbf{P}+\mathbf{p})\psi^*_{n_CL_CM_{L_C}}(\tfrac{m_3}{m_2+m_3}\mathbf{P}+\mathbf{p})\\ &\times\psi_{n_AL_AM_{L_A}}(\mathbf{P}+\mathbf{p})\mathcal{Y}^m_1(\mathbf{p}) \qquad (7)\end{aligned}$$

$$\begin{aligned}\mathcal{I}^{M_{L_A},m}_{M_{L_B},M_{L_C}}(-\mathbf{P},m_2,m_1,m_3) &= \sqrt{8E_AE_BE_C}\int d\mathbf{p}\,\psi^*_{n_BL_BM_{L_B}}(-\tfrac{m_3}{m_1+m_3}\mathbf{P}+\mathbf{p})\psi^*_{n_CL_CM_{L_C}}(-\tfrac{m_3}{m_2+m_3}\mathbf{P}+\mathbf{p})\\ &\times\psi_{n_AL_AM_{L_A}}(-\mathbf{P}+\mathbf{p})\mathcal{Y}^m_1(\mathbf{p}) \qquad (8)\end{aligned}$$

where $\mathbf{P_B} = -\mathbf{P_C} = \mathbf{P}, \mathbf{p} = \mathbf{p_3}$, and m_3 is the mass of the created quark. The spacial wavefunction one take is the simple harmonic oscillator (SHO) wavefunction. In momentum-space, the SHO wavefunction reads

$$\begin{aligned}\Psi_{nLM_L}(\mathbf{p}) &= (-1)^n(-i)^LR^{L+\frac{3}{2}}\sqrt{\frac{2n!}{\Gamma(n+L+\frac{3}{2})}}\\ &\times\exp\left(-\frac{R^2p^2}{2}\right)L^{L+\frac{1}{2}}_n(R^2p^2)\mathcal{Y}_{LM_L}(\mathbf{p}), \quad (9)\end{aligned}$$

here $\mathcal{Y}_{LM_L}(\mathbf{p})$ is the solid harmonic polynomial; R is the parameter of SHO wavefunction; $\mathbf{p}$ is the relative momentum between $q\bar{q}$ pair within one meson; $L^{L+\frac{1}{2}}_n(R^2p^2)$ is the Laguerre polynomial.

The decay width can be written as follows

$$\Gamma = \pi^2\frac{|\mathbf{P}|}{M^2_A(1+\delta_{BC})}\sum_{JL}\left|\mathcal{M}^{JL}\right|^2, \quad (10)$$

where $\mathcal{M}^{JL}$ is the partial wave amplitude, which is related to the helicity amplitude $\mathcal{M}^{M_{J_A}M_{J_B}M_{J_C}}$ via the Jacob-Wick formula [27]

$$\begin{aligned}\mathcal{M}^{JL}(A\to BC) &= \frac{\sqrt{2L+1}}{2J_A+1}\sum_{M_{J_B},M_{J_C}}\langle L0JM_{J_A}|J_AM_{J_A}\rangle\\ &\times\langle J_BM_{J_B}J_CM_{J_C}|JM_{J_A}\rangle\mathcal{M}^{M_{J_A}M_{J_B}M_{J_C}}(\mathbf{P}), \quad (11)\end{aligned}$$

where $\mathbf{J}=\mathbf{J}_B+\mathbf{J}_C$, $\mathbf{J}_A=\mathbf{J}_B+\mathbf{J}_C+\mathbf{L}$. Then the decay width in terms of the partial wave amplitude is taken as, where $|\mathbf{P}|=|\mathbf{P}_B|=|\mathbf{P}_C|$. According to the calculation of 2-body phase space, one can get

$$|\mathbf{P}|=\frac{\sqrt{[M_A^2-(M_B+M_C)^2][M_A^2-(M_B-M_C)^2]}}{2M_A},$$

where M_A, M_B, and M_C are the masses of the meson A, B, and C, respectively.

3 The masses of the mesons

To calculate the meson spectrum, a QCD-inspired model, constituent quark model, is used. The model incorporates the perturbative (one gluon exchange) and nonperturbative (color confinement and spontaneous breaking of chiral symmetry) properties of QCD. The constituent quark mass originates from the spontaneous breaking of chiral symmetry and consequently constituent quarks should interact through the exchange of Goldstone bosons [28], in addition to the one-gluon-exchange. To describe the hadron-hadron interaction, the chiral partner of pion, σ-meson, is also used. So the model Hamiltonian is

$$H=m_1+m_2+\frac{\mathbf{p}^2}{2\mu}+V^C+V^G+V^\chi+V^\sigma, \qquad (12)$$

$$V^C=\boldsymbol{\lambda}_1^c\cdot\boldsymbol{\lambda}_2^c\left[-a_c(1-e^{-\mu_c r})+\Delta\right]+V_{SO}^C$$

$$\begin{aligned}
V_{SO}^C &= -\boldsymbol{\lambda}_1^c\cdot\boldsymbol{\lambda}_2^c\frac{a_c\mu_c e^{-\mu_c r}}{4m_1^2m_2^2 r}\left[(m_1^2+m_2^2)(1-2a_s)\right. \\
&\quad \left.+4m_im_j(1-a_s)\right]\ \mathbf{S}\cdot\mathbf{L} \\
V^G &= V_C^G+V_{SO}^G+V_T^G \\
V_C^G &= \frac{\alpha_s}{4}\boldsymbol{\lambda}_1^c\cdot\boldsymbol{\lambda}_2^c\left\{\frac{1}{r}-\frac{\boldsymbol{\sigma}_1\cdot\boldsymbol{\sigma}_2}{6m_1m_2}\frac{e^{-r/r_0(\mu)}}{r\,r_0^2(\mu)}\right\} \\
V_{OGE}^{SO} &= -\frac{\alpha_s}{16}\frac{\boldsymbol{\lambda}_1^c\cdot\boldsymbol{\lambda}_2^c}{m_1^2m_2^2}\left[\frac{1}{r^3}-\frac{e^{-r/r_g(\mu)}}{r^3}\left(1+\frac{r}{r_g(\mu)}\right)\right] \\
&\quad \left[((m_1+m_2)^2+2m_1m_2)\,\mathbf{S}\cdot\mathbf{L}\right], \\
V_{OGE}^T &= -\frac{1}{16}\frac{\alpha_s}{m_1m_2}\boldsymbol{\lambda}_1^c\cdot\boldsymbol{\lambda}_2^c\left[\frac{1}{r^3}-\frac{e^{-r/r_g(\mu)}}{r}\right. \\
&\quad \left.\left(\frac{1}{r^2}+\frac{1}{3r_g^2(\mu)}+\frac{1}{r\,r_g(\mu)}\right)\right]S_{12}, \\
V_\chi &= \left(v_\pi^C+v_\pi^T\right)\sum_{a=1}^{3}\lambda_1^a\ \lambda_2^a+\left(v_K^C+v_K^T\right)\sum_{a=4}^{7}\lambda_1^a\ \lambda_2^a \\
&\quad +\left(v_\eta^C+v_\eta^T\right)\left(\lambda_1^8\ \lambda_2^8\cos\theta_P-\lambda_1^0\ \lambda_2^0\sin\theta_P\right) \\
v_\chi^C &= C_1\left[Y(m_\chi r)-\frac{\Lambda_\chi^3}{m_\chi^3}Y(\Lambda_\chi r)\right]\boldsymbol{\sigma}_1\cdot\boldsymbol{\sigma}_2, \\
v_\chi^T &= C_1\left[H(m_\chi r)-\frac{\Lambda_\chi^3}{m_\chi^3}H(\Lambda_\chi r)\right]S_{12} \\
C_1 &= \frac{g_{ch}^2}{4\pi}\frac{m_\chi^2}{12m_1m_2}\frac{\Lambda_\chi^2}{\Lambda_\chi^2-m_\chi^2}m_\chi, \quad \chi=\pi,K,\eta, \\
V_\sigma &= -C_2\left[Y(m_\sigma r)-\frac{\Lambda_\sigma}{m_\sigma}Y(\Lambda_\sigma r)\right]+V_\sigma^{SO}, \\
V_\sigma^{SO} &= -C_2\frac{m_\sigma^2}{2m_1m_2}\left[G(m_\sigma r)-\frac{\Lambda_\sigma^3}{m_\sigma^3}G(\Lambda_\sigma r)\right]\mathbf{S}\cdot\mathbf{L} \\
C_2 &= \frac{g_{ch}^2}{4\pi}\frac{\Lambda_\sigma^2}{\Lambda_\sigma^2-m_\sigma^2}m_\sigma
\end{aligned}$$

$$S_{12}=3(\boldsymbol{\sigma}_1\cdot\mathbf{r})(\boldsymbol{\sigma}_2\cdot\mathbf{r})-\boldsymbol{\sigma}_1\cdot\boldsymbol{\sigma}_2,$$

$$Y(x)=\frac{e^{-x}}{x}, \quad H(x)=\left(1+\frac{3}{x}+\frac{3}{x^2}\right)Y(x),$$

$$G(x)=\left(1+\frac{1}{x}\right)\frac{Y(x)}{x},$$

where $r=|\mathbf{r}_1-\mathbf{r}_2|$ and $\mathbf{p}=(\mathbf{p}_1-\mathbf{p}_2)/2$, $r_0(\mu)=\hat{r}_0/\mu$, $r_g(\mu)=\hat{r}_g/\mu$. Other symbols have their usual meanings. The effective running coupling constant is given by

$$\alpha_s(\mu)=\frac{\alpha_0}{\ln\left(\frac{\mu^2+\mu_0^2}{\Lambda_0^2}\right)}, \qquad (13)$$

where μ is the reduced mass of the $q\bar{q}$ system. The chiral coupling constant g_{ch} is determined from the πNN coupling constant through

$$\frac{g_{ch}^2}{4\pi}=\left(\frac{3}{5}\right)^2\frac{g_{\pi NN}^2}{4\pi}\frac{m_{u,d}^2}{m_N^2}. \qquad (14)$$

The meson spectrum is obtained by solving the Schrödinger equation,

$$\begin{aligned} H\Psi &= E\Psi, & (15)\\ \Psi &= \left[\psi_{nLM_L}\chi_{SM_S}\right]_{JM_J}\chi_c\chi_f, & (16) \end{aligned}$$

Table 1. Model parameters. The masses of mesons π, K, η take the experimental values.

$m_{u,d}$	m_s	a_c	μ_c	Δ	a_s
MeV	MeV	MeV	fm^{-1}	MeV	-
313	555	430	0.7	181.10	0.777
α_0	Λ_0	μ_0	$\hat{r}_0$	$\hat{r}_g$	
-	fm^{-1}	MeV	MeV fm	MeV fm	
2.118	0.113	36.976	28.170	34.500	
Λ_π	Λ_σ	Λ_K	Λ_η	$g^2_{ch}/4\pi$	θ_P
fm^{-1}	fm^{-1}	fm^{-1}	fm^{-1}	-	^{o}
4.20	4.20	5.20	5.20	0.54	-15

where $\chi_{SM_S}, \chi_c, \chi_f$ are spin, color and flavor wavefunctions of the meson, respectively and can be constructed through the symmetry. The spatial wavefunction $\psi_{nLM_L} = R_{nL}(r)Y_{LM_L}(\Omega)$ is obtained by solving the second-order differential equation. The efficient numerical method: Numerov method [29] is used here. The model parameters, which are listed in Table 1, are fixed by fitting the experimental data of meson spectrum. Parts of the obtained meson spectrum are shown in Tables 2 and 3. The detailed results can be found in Ref. [20]. To calculate the strong decay of mesons analytically in $^3P_0$ model, the obtained radial part of the spacial wavefunction $R_{nL}(r)$ is fitted by the simple harmonic oscillator (SHO),

$$\begin{aligned} R_{nL}(r) &= \beta^{(L+\frac{3}{2})}\sqrt{\frac{2n!}{\Gamma(n+L+\frac{3}{2})}}\exp\left(-\frac{\beta^2r^2}{2}\right) \\ & \quad r^L L_n^{L+\frac{1}{2}}(\beta^2r^2). \qquad (17) \end{aligned}$$

The fitted values of parameter β are also listed in Table 3.

Table 2. The mass of $I=1,\frac{1}{2}$ mesons and the values of fitted β.

$n^{2S+1}L_J$	states	Isospin	Mass (MeV)	β (fm^{-1})	R (GeV^{-1})
1^1S_0	π	1	139	2.308	2.196
2^1S_0	$\pi(1300)$	1	1288	1.434	3.534
1^3S_1	ρ	1	772	1.438	3.522
2^3S_1	$\rho(1450)$	1	1478	1.096	4.624
1^1P_1	$b_1(1235)$	1	1234	1.243	4.077
1^3P_0	$a_0(980)$	1	984	1.473	3.440
2^3P_0	$a_0(1450)$	1	1587	1.125	4.505
1^3P_1	$a_1(1260)$	1	1205	1.300	3.898
1^3P_2	$a_2(1320)$	1	1327	1.106	4.582
1^3P_2	$a_2(1700)$	1	1732	0.890	5.694
1^1S_0	K	1/2	496	2.313	2.191
2^1S_0	$K(1460)$	1/2	1472	1.545	3.280
1^3S_1	$K^*(892)$	1/2	910	1.629	3.111
2^3S_1	$K(1630)$	1/2	1620	1.262	4.016
1^1P_1	$K_1(1400)$	1/2	1414	1.371	3.696
1^3P_0	$K_0^*(1430)$	1/2	1213	1.572	3.224
2^3P_0	$K_0^*(1950)$	1/2	1768	1.243	4.077
1^3P_1	$K_1(273)$	1/2	1352	1.435	3.531
1^3P_2	$K_2^*(1430)$	1/2	1450	1.572	3.224
1^3D_1	$K_1(1680)$	1/2	1698	1.205	4.206

For $I=0$ states, there are two types of them, one is composed of u,d-quark and $\bar{u},\bar{d}$-antiquark, another is composed of s-quark and $\bar{s}$-antiquark. They are mixed in the flavor SU(3) symmetry to form flavor singlet and octet. However, flavor SU(3) is broken. In experiments, we have η and η' instead of η_1 and η_8 for pseudoscalar. In the present calculation, flavor SU(3) symmetry is not used, so we have flavor wavefunctions X_n and X_s. As a consequence of K-meson exchange, they are mixed. To obtain the masses of $I=0$ states, the following procedure is taken. First, solving the Schrödinger equation for X_n and X_s separately (K-meson exchange is not employed). Secondly, by using the wavefunctions Ψ_n and Ψ_s obtained in the first step and taking into account of K-meson exchange, the eigen-energies and eigen-states can be obtained by diagonalizing the Hamiltonian matrix

$$\begin{pmatrix} H_{nn} & H_{ns} \\ H_{sn} & H_{ss} \end{pmatrix}\begin{pmatrix} C_n \\ C_s \end{pmatrix} = E\begin{pmatrix} C_n \\ C_s \end{pmatrix}. \qquad (18)$$

where $H_{nn} = \langle\Psi_n|H|\Psi_n\rangle$, $H_{ns} = \langle\Psi_n|V_K|\Psi_s\rangle = H_{sn}$ and $H_{ss} = \langle\Psi_s|H|\Psi_s\rangle$. The eigen-state is $|\Psi\rangle = C_n|\Psi_n\rangle + C_s|\Psi_s\rangle$. The obtained eigen-energies and eigen-states are shown in Table 3. From Table 3, one finds that $\eta(1760), X(1835), X(2120), X(2370)$ may be interpreted as $\eta'(2^1S_0), \eta(4^1S_0), \eta'(3^1S_0)$ and $\eta'(4^1S_0)$ respectively by comparing the theoretical masses with the experimental data. To check these assignments, the decay properties of the states should be studied, which is discussed in the next section.

Table 3. The masses of $I=0$ mesons and the value of fitted β ($\beta=C_n^2\beta_n+C_s^2\beta_s$).

$(nL)J^{PC}$	states	Mass (MeV)	C_n	C_s	$\beta(\text{fm}^{-1})$	$R(\text{GeV}^{-1})$
1^1S_0	η	572	8.6564×10^{-1}	-5.0066×10^{-1}	1.732693	2.924
1^1S_0	$\eta'(958)$	956	5.0066×10^{-1}	8.6564×10^{-1}	2.064307	2.455
2^1S_0	$\eta(1295)$	1290	9.6360×10^{-1}	-2.67323×10^{-1}	1.183-1.666	3.041-4.284
2^1S_0	$\eta'(1760)$	1795	2.6732×10^{-1}	9.6360×10^{-1}	1.183-1.666	3.041-4.284
3^1S_0	$\eta(3S)$	1563	9.9350×10^{-1}	-1.1380×10^{-1}	0.929-1.360	3.726-5.455
3^1S_0	$\eta'(3S)$	2276	1.1380×10^{-1}	9.9350×10^{-1}	0.929-1.360	3.726-5.455
4^1S_0	$\eta(4S)$	1807	9.9935×10^{-1}	-3.5928×10^{-2}	0.6725-1.0995	4.607-7.530
4^1S_0	$\eta'(4S)$	2390	3.5928×10^{-2}	9.9935×10^{-1}	0.6725-1.0995	4.607-7.530
1^3S_1	$\omega(782)$	691	9.9499×10^{-1}	9.9967×10^{-2}	1.547	3.276
1^3S_1	$\phi(1020)$	1020	-9.9967×10^{-2}	9.9499×10^{-1}	1.918	2.642
2^3S_1	$\omega(1420)$	1444	9.9852×10^{-1}	5.4331×10^{-2}	1.163	4.357
2^3S_1	$\phi(1680)$	1726	-5.4331×10^{-2}	9.9852×10^{-1}	1.506	3.365
1^1P_1	$h_1(1170)$	1257	1.0	0	1.202	4.216
1^1P_1	h_1'	1511	0	1.0	1.581	3.205
1^3P_2	$f_2(1270)$	1311	1.0	0	1.112	4.557
1^3P_2	$f_2'(1525)$	1556	0	1.0	1.496	3.387

4 The strong decay of the candidates for $\eta(1760), X(1835), X(2120), X(2370)$

η, η' and their radial excitations have the same quantum numbers $IJ^{PC}=00^{-+}$. According to the $^3P_0$ model discussed above, the isospins of mesons B and C can takes the values $I=0,1/2$, or 1 with the condition $\mathbf{I}_B+\mathbf{I}_C=\mathbf{I}_A$. If not forbidden kinetically, the allowed decay modes of $\eta(\eta')$ family are listed in Table 4.

Table 4. Allowed decay modes and the amplitudes of the radial excited states of η and η'. For X_n decay, $\phi_f=\sqrt{\frac{1}{2}},\ \sqrt{\frac{1}{3}},\ \sqrt{\frac{1}{6}},\ 0$ for $I_B=I_C=1,\ 1/2,\ 0(X_n),\ 0(X_s)$ and for X_s decay, $\phi_f=0,\ \sqrt{\frac{2}{3}},\ 0,\ \sqrt{\frac{1}{3}}$ for $I_B=I_C=1,\ 1/2,\ 0(X_n),\ 0(X_s)$.

$X \to {}^1S_0 + {}^3P_0$	$\pi a_0(980),\ \pi a_0(1450),\ \pi(1300)a_0(980),$ $KK_0^*(1430),\ KK_0^*(1950)$	$M^{JL}=M^{00}=M^{000}$ $M^{000}=\sqrt{\frac{1}{36}}(I_{0,0}^{-1,-1}+I_{0,0}^{0,0}+I_{0,0}^{1,1})\phi_f$
$X \to {}^1S_0 + {}^3P_2$	$\pi a_2(1320),\ \pi a_2(1700),\ KK_2^*(1430),\ \eta f_2(1270),$ $\eta' f_2(1270),\ \eta f_2'(1525)$	$M^{JL}=M^{22}=M^{000}$ $M^{000}=\sqrt{\frac{1}{72}}(I_{0,0}^{-1,-1}-2M^{0-11})\phi_f$
$X \to {}^1S_0 + {}^3S_1$	$KK^*,\ KK^*(1410),\ K(1460)K^*$	$M^{JL}=M^{11}=-M^{000}$ $M^{000}=-\sqrt{\frac{1}{12}}I_{0,0}^{0,0}\phi_f$
$X \to {}^1S_0 + {}^3D_1$	$KK^*(1680)$	$M^{JL}=M^{11}=-M^{000}$ $M^{000}=(\sqrt{\frac{1}{40}}I_{0,0}^{-1,-1}+\sqrt{\frac{1}{30}}I_{0,0}^{0,0}+\sqrt{\frac{1}{40}}I_{0,0}^{1,1})\phi_f$
$X \to {}^3S_1 + {}^3P_1$	$\rho a_1(1640),\ \rho a_1(1260),\ K^*K_1(1273),\ \omega f_1(1285)$	$M^{JL}=M^{00}+M^{22}$ $M^{00}=\sqrt{\frac{1}{3}}(M^{0-11}-M^{000}+M^{01-1})$ $M^{22}=\sqrt{\frac{1}{6}}(M^{0-11}+2M^{000}+M^{01-1})$ $M^{0-11}=-\sqrt{\frac{1}{24}}(I_{0,0}^{0,0}+I_{0,0}^{1,1})\phi_f$ $M^{000}=\sqrt{\frac{1}{24}}(I_{0,0}^{-1,-1}+I_{0,0}^{1,1})\phi_f$ $M^{01-1}=-\sqrt{\frac{1}{24}}(I_{0,0}^{0,0}+I_{0,0}^{1,1})\phi_f$
$X \to {}^3S_1 + {}^3S_1$	$\rho\rho,\ \rho\rho(1450),\ \omega\omega,\ \omega\omega(1420),$ $K^*K^*,\ K^*K^*(1410),\ \phi\phi$	$M^{JL}=M^{11}=\sqrt{\frac{1}{2}}(M^{0-11}-M^{01-1})$ $M^{0-11}=\sqrt{\frac{1}{12}}(I_{0,0}^{0,0})\phi_f,\ M^{0-11}=-\sqrt{\frac{1}{12}}(I_{0,0}^{0,0})\phi_f$
$X \to {}^3S_1 + {}^1P_1$	$\rho b_1(1235),\ K^*K_1(1400),\ \omega h_1(1170)$	$M^{JL}=M^{00}+M^{22}$ $M^{00}=\sqrt{\frac{1}{3}}(M^{0-11}-M^{000}+M^{01-1})$ $M^{22}=\sqrt{\frac{1}{6}}(M^{0-11}+2M^{000}+M^{01-1})$ $M^{0-11}=\sqrt{\frac{1}{12}}I_{0,0}^{1,1}\phi_f,\quad M^{000}=-\sqrt{\frac{1}{12}}I_{0,0}^{1,1}\phi_f$ $M^{01-1}=\sqrt{\frac{1}{12}}I_{0,0}^{-1,-1}\phi_f$
$X \to {}^3S_1 + {}^3P_2$	$\rho a_2(1320),\ K^*K_2^*(1430)$	$M^{JL}=M^{22}=-\sqrt{\frac{1}{2}}(M^{0-11}-M^{01-1})$ $M^{0-11}=\sqrt{\frac{1}{24}}(I_{0,0}^{0,0}-I_{0,0}^{1,1})\phi_f$ $M^{01-1}=\sqrt{\frac{1}{24}}(I_{0,0}^{-1,-1}-I_{0,0}^{1,1})\phi_f$

All the possible decay modes of $\eta(\eta')$ family are shown in Table 4. To calculate the strong decay widths of mesons, the strength of the quark pair creation from the vacuum, γ, has to be fixed. It is obtained by fitting the experimental values of the strong decay widths of light and charmed mesons, charmonium and baryons. In the present work, $\gamma = 6.95$, which is adopted by many researches [30–32], is taken for the non-strange quark pair creation, and the strength of $s\bar{s}$ creation satisfies γ_s=$\gamma/3$ [33].

4.1 $\eta'(2^1S_0)$

The experimental evidence for $\eta(1760)$ is controversial. There are large differences between the observations of MARK III, DM2 and BES collaborations[1, 3–6]. In our calculation, the mass of $\eta'(2^1S_1)$ is 1795 MeV, which is close to the experimental mass of $\eta(1760)$. So we take it as the candidate of $\eta(1760)$. In Fig.2, we show the dependence of the partial widths of the strong decay of the $\eta'(2^1S_0)$ on the R_A. Taking R_A =3.0-4.3 GeV^{-1} discussed above, the total width ranges from 256 to 404 MeV, which is much larger than the results given by Mark III and DM2 collaboration, but falls in the range of the BES experimental data. In this range, $\eta'(2^1S_0)$ have a sizable branching ration into $\pi a_0(980)$, $\pi a_2(1320)$, $\rho\rho$, and KK^*. But the partial width to $\omega\omega$ is rather small. If the BES results are reliable, the assignment of $\eta(1760)$ to $\eta'(2^1S_0)$ is disfavored in the present calculation. In Ref. [19], $\eta(1760)$ is taken as $\eta(3S)$, the total decay width is between 60-100 MeV, which falls in the range of DM2's results, but is far below BES's results.



Fig. 2. The possible strong decay of the $\eta'(2^1S_0)$

4.2 $\eta(4^1S_0)$

$X(1835)$ was first observed by BESII in the $\pi^+\pi^-\eta'$ invariant-mass spectrum in the decay channel $J/\psi \to \gamma\pi^+\pi^-\eta'$ with a statistical significance of 7.7 σ[1]. BESIII confirmed it in the same process with statistical significance larger than 20σ [2]. In the present calculation, the mass of $\eta(4^1S_0)$=1807 MeV is close to the mass of $X(1835)$, so the assignment of $X(1835)$ to $\eta(4^1S_0)$ is possible, which is different from the assignment of Ref. [19], $\eta'(3S)$. In Fig. 2, the dependence of the partial widths of the strong decay of the $\eta(4^1S_0)$ on the R_A is shown. From the mass calculation, R_A =4.6-7.5 GeV^{-1} is obtained. In this range, the total width ranges from 54 to 692 MeV, which falls in the range of the BES experimental data, and the main decay modes are $\pi a_0(980)$ and $\pi a_0(1450)$. We suggest experimental search for $X(1835)$ in these modes to make sure whether it is $\eta(4^1S_0)$ assignment.



Fig. 3. The possible strong decay of the $\eta(4^1S_0)$



Fig. 4. The possible strong decay of the $\eta'(3^1S_0)$

4.3 $\eta'(3^1S_0)$ and $\eta'(4^1S_0)$

Besides confirmed the existence of $X(1835)$ in the $\pi^+\pi^-\eta'$ invariant-mass spectrum in the process $J/\psi \to \eta'\pi^+\pi^-$, other two states $X(2120)$ and $X(2370)$ are observed by BESIII with statistical significance larger than 7.2σ and 6.4 σ, respectively. By comparing the masses of the $\eta(\eta')$ family, it is possible to take $\eta'(3^1S_0)$ and $\eta'(4^1S_0)$ as the candidates of $X(2120)$ and $X(2370)$. Because of their large masses, many strong decays modes are allowed. In Figs. 4 and 5, the partial widths of their strong decays are shown. For $\eta'(3^1S_0)$ with R_A =3.7-5.6 GeV^{-1} and for $\eta'(4^1S_0)$ with R_A =4.6-7.5 GeV^{-1}, the decay widths are much higher than the experimental data of BESIII. Because both X_n and X_s have contributions to the state $n\bar{s}s\bar{n}$, the partial width of the strong decay to two isospin I=$\frac{1}{2}$ mesons is generally much larger that to two isospin 1 or 0 mesons. $\eta'(3^1S_0)$ have large partial to KK_0^* and KK^*. And the main decay modes of $\eta'(4^1S_0)$ are KK^*, $KK_1(1352)$, $KK_0^*(1430)$, $KK_0^*(1950)$.

If we describe $X(2120)$ and $X(2370)$ as $\eta'(3^1S_0)$ and $\eta'(4^1S_0)$ respectively with parameters in this work , it is not appropriate obviously.



Fig. 5. The possible strong decay of the $\eta'(4^1S_0)$

5 summary and discussions

By using chiral quark model, the mass spectrum of $\eta(\eta')$ family are calculated, where the mixing between $(u\bar{u}+d\bar{d})/\sqrt(2)$ and $s\bar{s}$ is determined by system dynamics, K-meson exchange. Based on the mass spectrum, the possible candidates of four $J^{PC}I^G=0^{-+}0^+$ mesons, $\eta(1760)$, $X(1835)$, $X(2120)$ and $X(2370)$ are assigned to $\eta'(2^1S_0)$, $\eta(4^1S_0)$, $\eta'(3^1S_0)$, $\eta'(4^1S_0)$. Furthermore, all kinematically allowed two-body strong decays of them can calculated in the framework of the $^3P_0$ model. The wavefunctions needed in the calculation are obtained from the mass calculation. To simplify the calculation, the SHO wavefunctions are used to mimic the real wavefunctions.

The decay widths turn out to be strongly dependent on the SHO wave function scale parameter β. For $\eta(1760)$, the width is larger than the result of [6] and is compatible with the results of BES observation [1] in the R_A range. However the partial width to $\omega\omega$ is too small, which it is incompatible with experimental date [1, 3, 6]. So the assignment of $eta(1760)$ to $\eta(2S)$ is disfavored in the present calculation. For the state $X(1835)$, the calculated decay width is consistent with experiment data, and $\pi a_0(980)$ and $\pi a_0(1450)$ are the main decay modes. To justify this assignment, the experimental investigation of the $\pi a_0(980)$ and $\pi a_0(1450)$ decay modes of $X(1835)$ is needed. Sine $X(1835)$ is around the threshold of $p\bar{p}$, it may be the the mixture of $q\bar{q}$ and baryonium. Further study of the state $X(1835)$ by taking into account of the mixture is essential to understand the nature of the state.

$X(2120)$ and $X(2370)$ are assigned to $\eta'(3^1S_0)$ and $\eta'(4^1S_0)$ respectively. Since they have larger masses, many strong decays modes are allowed and have a large phase space to some modes. The total decay widths are much higher than the experimental values. The large decay width may de due to the overestimated value of γ. To exclude the impact of parameters, the branching ratio is better to justify the assignment. More experimental data are needed. Since the lattice QCD predicts the 0^{-+} glueball is about 2.3∼2.6 GeV, which is around the masses of $X(2120)$ and $X(2370)$, the study with the mixture of $q\bar{q}$, glueball and other configurations are necessary to understand the nature of $X(2120)$ and $X(2370)$ states.

References

1 Ablikim M et al. [BES Collaboration], Phys. Rev. Lett., 2005, **95**, 262001:1-5
2 Ablikim M et al. [BES Collaboration], Phys. Rev. Lett., 2011, **106**, 072002:1-5
3 Baltrusaitis R M et al. [MARKIII Collaboration], Phys. Rev. Lett., 1985, **55**, 1723-1726
4 Baltrusaitis R M et al. [MARKIII Collaboration], Phys. Rev. D, 1986, **33**, 1222-1232
5 Bisello D et al. [DM2 Collaboration], Phys. Lett. B, 1987, **192**, 239-244
6 Bisello D et al. (DM2 Collaboration), Phys. Rev. D, 1989, **39**, 701-712
7 Ablikim M et al. [BES Collaboration], Phys. Rev. D, 2006, **73**, 112007:1-9
8 Klempt E and Zaitsev A, Phys. Rep., 2007, **454**, 1-202 and refrences therein.
9 Ding G J and Yan M L, Phy Rev. C, 2005, **72**, 015208:1-8; Ding G J and Yan M L, Eur. Phys. J. A, 2006, **28**, 351-360
10 Dedonder J P, Loiseau B, El-Bennich B and Wycech S, Phys. Rev. C, 2009, **80**, 045207:1-6
11 Liu C, Eur. Phys. J. C, 2008, **53**, 413-419
12 Wang Z G and Wan S L, J. Phys. G, 2007, **34**, 505-512
13 Kochelev N and Min D P, Phys. Lett. B, 2006, **633**, 283-288; Phys. Rev. D, 2005, **72**, 097502:1-3
14 He X G, Li X Q, Liu X and Ma J P, Eur. Phys. J. C, 2006, **49**, 731-736
15 Li B A, Phys. Rev. D, 2006, **74**, 034019:1-8
16 Hao G, Qiao C F and Zhang A L, Phys. Lett. B, 2006 **642**, 53-61
17 Huang T and Zhu S L, Phys. Rev. D, 2006, **73**, 014023:1-4
18 D. M. Li and B. Ma, Phys. Rev. D 77, 074004:1-7 (2008).
19 Yu J S, Sun Z F, Liu X and Zhao Q, Phys. Rev. D, 2011, **83**, 114007:1-15
20 Vijande J, Fernández F and Valcarce A, J. Phys. G, 2005, **31**, 481-506
21 Li X Q and Page P R, Eur. Phys. J. C, 1998, **1**, 579-583
22 Wu N, Yuan T N and Zheng Z P, Chin. Phys. C, 2001, **10**, 611-612
23 Micu L, Nucl. Phys. B, 1969, **10**, 521-526
24 Yaouanc A Le, Oliver L, Pene O and Raynal J C, Phys. Rev. D, 1973, **8**, 2223-2234; 1974, **9**, 1415-1419; 1975, **11**, 1272-1286
25 Roberts W and Silvestr-Brac B, Few-Body Syst., 1992, **11**, 171-193
26 Ackleh E S, Barnes T and Swanson E S, Phys. Rev. D, 1996, **54**, 6811-6829
27 Jacob M and Wick G C, Ann. Phys. (N.Y.), 1959, **7**, 404-428
28 Manohar A and Georgi H, Nucl. Phys. B, 1984, **234**, 189-212
29 Koonin S E and Meredith D C. Computational Physics. New York: Addison-Wesley, 1990. 56—57
30 Lu J, Deng W Z, Chen X L and Zhu S L, Phys. Rev. D, 2006, **73**, 054012:1-6; Zhang B, Liu X, Deng W Z and Zhu S L, Eur. Phys. J. C, 2007, **50** 617-628; Chen C, Chen X L, Liu X, Deng W Z and Zhu S L, Phys. Rev. D, 2007, **75**, 094017:1-13
31 Blundell H G and Godfrey S, Phys. Rev. D, 1996, **53**, 3700-3711; Phys. Rev. D, 1996, **53**, 3712-3722
32 Yang Y C, Xia Z R and Ping J L, Phys. Rev. D, 2010, **81**, 094003:1-9
33 Yaouanc A Le, Oliver L, Pene O and Raynal J C, Phys. Lett. B, 1977, **72**, 57-61